Table 10 (page 1 of 2). Infant, neonatal, postneonatal, fetal, and perinatal mortality rates, by detailed race and Hispanic origin of mother: United States, selected years 1983–2014

Excel and PDF versions (with more data years and standard errors when available): http://www.cdc.gov/nchs/hus/contents2016.htm#010.

[Data are based on linked birth and death certificates for infants and fetal death records]

Maternal race and Hispanic origin	1983 ¹	1985 ¹	1990 ¹	1995²	2000 ²	2005 ²	2010 ²	2013 ²	2014 ²	
	Infant ³ deaths per 1,000 live births									
All mothers	10.9	10.4	8.9	7.6	6.9	6.9	6.1	6.0	5.8	
White	9.3 19.2 15.2 8.3	8.9 18.6 13.1 7.8	7.3 16.9 13.1 6.6	6.3 14.6 9.0 5.3	5.7 13.5 8.3 4.9	5.7 13.3 8.1 4.9	5.2 11.2 8.3 4.3	5.1 10.8 7.6 4.1	4.9 10.7 7.6 3.9	
Hispanic or Latina 5.6	9.5 9.1 12.9 7.5 8.5 10.6	8.8 8.5 11.2 8.5 8.0 9.5	7.5 7.2 9.9 7.2 6.8 8.0	6.3 6.0 8.9 5.3 5.5 7.4	5.6 5.4 8.2 4.6 4.6 6.9	5.6 5.5 8.3 4.4 4.7 6.4	5.3 5.1 7.1 3.8 4.4 6.1	5.0 4.9 5.9 3.0 4.3 5.9	5.0 4.8 7.2 3.9 4.3 5.4	
White	9.2 19.1	8.6 18.3	7.2 16.9	6.3 14.7	13.6	5.8 13.6	5.2 11.5	11.1	10.9	
	Neonatal ³ deaths per 1,000 live births									
All mothers	7.1	6.8	5.7	4.9	4.6	4.5	4.0	4.0	3.9	
White	6.1 12.5 7.5 5.2	5.8 12.3 6.1 4.8	4.6 11.1 6.1 3.9	4.1 9.6 4.0 3.4	3.8 9.1 4.4 3.4	3.8 8.9 4.0 3.4	3.5 7.3 4.3 3.0	3.4 7.3 4.1 3.0	3.3 7.3 4.1 2.8	
Hispanic or Latina 5,6. Mexican	6.2 5.9 8.7 *5.0 5.8 6.4	5.7 5.4 7.6 6.2 5.6 5.6	4.8 4.5 6.9 5.3 4.4 5.0	4.1 3.9 6.1 *3.6 3.7 4.8	3.8 3.6 5.8 *3.2 3.3 4.6	3.9 3.8 5.9 *3.1 3.2 4.3	3.6 3.5 4.8 *2.9 3.0 4.0	3.6 3.5 4.2 *2.3 3.1 4.0	3.5 3.5 5.0 2.6 3.1 3.7	
Not Hispanic or Latina: 6 White	5.9 12.0	5.6 11.9	4.5 11.0	4.0 9.6	3.8 9.2	3.7 9.1	3.4 7.5	3.3 7.5	3.2 7.4	
	Postneonatal ³ deaths per 1,000 live births									
All mothers	3.8	3.6	3.2	2.6	2.3	2.3	2.1	1.9	1.9	
White	3.2 6.7 7.7 3.1	3.1 6.3 7.0 2.9	2.7 5.9 7.0 2.7	2.2 5.0 5.1 1.9	1.9 4.3 3.9 1.4	2.0 4.3 4.0 1.5	1.8 3.9 4.0 1.3	1.6 3.5 3.5 1.1	1.6 3.4 3.5 1.0	
Hispanic or Latina 5,6	3.3 3.2 4.2 *2.5 2.6 4.2	3.2 3.2 3.5 *2.3 2.4 3.9	2.7 2.7 3.0 *1.9 2.4 3.0	2.1 2.1 2.8 *1.7 1.9 2.6	1.8 1.8 2.4 * 1.4 2.3	1.8 1.7 2.4 *1.4 1.5 2.1	1.7 1.6 2.3 * 1.4 2.1	1.5 1.4 1.7 * 1.2 1.9	1.5 1.4 2.2 *1.3 1.1 1.7	
Not Hispanic or Latina: 6 White	3.2 7.0	3.0 6.4	2.7 5.9	2.2 5.0	1.9 4.4	2.1 4.5	1.8 4.0	1.7 3.7	1.7 3.5	

See footnotes at end of table.

Health, United States, 2016 Trend Tables 107

Table 10 (page 2 of 2). Infant, neonatal, postneonatal, fetal, and perinatal mortality rates, by detailed race and Hispanic origin of mother: United States, selected years 1983–2014

Excel and PDF versions (with more data years and standard errors when available): http://www.cdc.gov/nchs/hus/contents2016.htm#010.

[Data are based on linked birth and death certificates for infants and fetal death records]

Maternal race and Hispanic origin	1983	1985	1990	1995	2000 ⁷	2005	2010	2013	2014	
тіізрапіс опуш	1900	1900	1990		2000	2005	2010	2013	2014	
	Fetal 8,9 deaths per 1,000 live births plus fetal deaths									
All mothers				7.0	6.6	6.2	6.0	6.0	6.0	
Hispanic or Latina⁵					5.8	5.4	5.2	5.2	5.1	
White					5.3	4.8	4.8	4.9	4.9	
Black or African American					12.0	11.1	10.8	10.5	10.7	
	Late fetal 8,10 deaths per 1,000 live births plus late fetal deaths									
All mothers				3.6	3.3	3.0	3.0	3.0	2.8	
Hispanic or Latina ⁵					3.1	2.8	2.6	2.7	2.5	
White					2.8	2.4	2.5	2.6	2.4	
Black or African American					5.2	4.8	4.7	4.7	4.6	
	Perinatal 8,11 deaths per 1,000 live births plus late fetal deaths									
All mothers				7.6	7.0	6.6	6.2	6.2	6.0	
Hispanic or Latina⁵					6.1	5.9	5.5	5.6	5.4	
White					5.7	5.4	5.1	5.3	5.0	
Black or African American					12.6	12.2	10.6	10.7	10.6	

^{*} Estimates are considered unreliable. Rates preceded by an asterisk are based on fewer than 50 deaths in the numerator. Rates not shown are based on fewer than 20 deaths in the numerator.

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander include persons of Hispanic and non-Hispanic origin. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. National linked files do not exist for 1992–1994. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: http://www.cdc.gov/nchs/hus.htm.

SOURCE: NCHS, National Vital Statistics System, public-use Linked Birth/Infant Death Data Set, public-use Fetal Death File, public-use Birth File. National Center for Health Statistics. Mathews TJ, Driscoll AK. Trends in infant mortality in the United States, 2005–2014. NCHS data brief, no 279. Hyattsville, MD: NCHS; 2017. Available from: https://www.cdc.gov/nchs/products/databriefs.htm and National Center for Health Statistics. 2014 fetal death data set and user's guide. Hyattsville, MD: 2016. Available from: https://www.cdc.gov/nchs/data_access/vitalstatsonline.htm. See Appendix I, National Vital Statistics System (NVSS).

108 Trend Tables Health, United States, 2016

^{- - -} Data not available.

¹ Rates based on unweighted birth cohort data.

² Rates based on a period file using weighted data. See Appendix I, National Vital Statistics System (NVSS), Linked Birth/Infant Death Data Set.

³ Infant (under 1 year of age), neonatal (under 28 days), and postneonatal (28 days-11 months).

⁴Estimates are not available for Asian or Pacific Islander subgroups because not all states have adopted the 2003 revision of the U.S. Standard Certificate of Live Birth. See Appendix II, Race.

⁵ Persons of Hispanic origin may be of any race.

⁶ Prior to 1995, data are shown only for states with an Hispanic-origin item on their birth certificates. See Appendix II, Hispanic origin.

⁷Rates for 1999–2004 (shown in spreadsheet version) exclude data from Oklahoma, which did not report Hispanic origin on the fetal death report in those years.

⁸ Starting with 2014 data, the obstetric estimate of gestation at delivery replaced the gestational age measure based on the date of the last normal menses, which was used for prior years. For more information on the impact of this change, see Appendix I, National Vital Statistic System (NVSS).

⁹ Number of fetal deaths of 20 weeks or more gestation per 1,000 live births plus fetal deaths.

¹⁰ Number of fetal deaths of 28 weeks or more gestation (late fetal deaths) per 1,000 live births plus late fetal deaths.

¹¹ Number of late fetal deaths plus infant deaths within 7 days of birth per 1,000 live births plus late fetal deaths.